

Mastermectin Pour-on

In cattle: For the treatment of infections with the following parasites Gastro-intestinal worms

- Haemonchus placei (adult and fourth stage larvae) Ostertagia ostertagi (adult and fourth stage larvae, including inhibited larvae) Trichostrongylus axei (adult and fourth stage larvae) Trichostrongylus colubriformis (adult and fourth stage larvae) Cooperia punctata (adult only) Cooperia oncophora (adult only) Strongyloides papillosus (adult only) Oesophagostomum radiatum, (adult and fourth stage larvae).
- Lungworm (adult and fourth stage larvae) – Dictyocaulus viviparus.
- Warbles (parasitic stages) – Hypoderma bovis, Hypoderma lineatum.
- Mange mites – Sarcoptes scabiei var. bovis. The product may also be used to reduce infection of the mange mite Chorioptes bovis, but complete elimination may not occur.
- Sucking and biting lice – Linognathus vituli, Haematopinus eurysternus, Bovicola (Damalinia) bovis.
- Mastermectin 0.5% Cattle Pour-On has persistent activity against infections acquired with Trichostrongylus axei and Cooperia spp. up to 14 days after treatment, but only in the case of group treatment; Ostertagia ostertagi and Oesophagostomum radiatum up to 21 days after treatment; Dictyocaulus viviparus up to 28 days after treatment. It also has persistent activity against horn flies (Haematobia irritans) for up to 28 days after treatment; partial efficacy against Haematobia irritans may last for up to 35 days post application. Health Products Regulatory Authority.

Benefits of using Ivermectin Pour-on

- Ease of use.
- Less stress on Animal and Farmer.
- Controls key internal parasites.
- Controls key external parasites.
- Good length of activity and efficacy.
- Wide margin of safety when used as directed read SPC Sheet.
- Rainfast.
- Proven & Registered by National Regulatory Authority - HPRA.